

Early Journal Content on JSTOR, Free to Anyone in the World


This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.



BP make equal angles with the radius drawn to P.

This is known as Alhazen's Problem and does not admit of a solution by ruler and compasses only. However, by the use of an hyperbola an approximate solution can be effected.

Construction. Let O be the center of the circle, a its radius. Take C and D so that $AO.OC = a^2 = BO.OD$.

Now, the locus of the vertices of the triangles whose base is CD and whose base angles have a constant difference (OCD-ODC) is well known to be a hyperbola. This will cut the circle in four points, of which let P be one. This is the point required.

Proof. $\angle CDP - \angle DCP = \angle OCD - \angle ODC$ (by construction). Transposing and adding,

 $\therefore \angle OCP = \angle ODP...(1)$.

Also in the triangles AOP, POC, we have AO:OP=OP:OC (by construction).

- $\therefore \angle APO = \angle OCP$. Similarly, $\angle BPO = \angle ODP$ by (1).
- $\therefore \angle APO = \angle BPO$, and $\therefore \angle APR = \angle BPR$. Q. E. F. Q. E. D.

D

350. Proposed by G. B. M. ZERR, A. M., Ph. D., Philadelphia, Pa.

Given the quadrilateral AB=a=225, BC=b=153, CD=c=207, DA=d=135, AC=e=240. Find the side of the square inscribed in this quadrilateral having a corner in each side.

Solution by the PROPOSER.

Let ABCD be the given quadrilateral, EFGH the inscribed square, RPQS the circumscribed rectangle having its sides parallel to the sides of the square. Draw AIJ, BUT, CNL, DVM perpendicular, respectively, to EF and HG, FG and HE, HG and EF, HE and GF.

Let AB=a, BC=b, CD=c, DA=d, AC=e, BD=f, O the intersection of AC,

BD, $\angle COB = \beta$, $\angle ACL = \theta = \angle CAJ$, $\angle BDM = \phi = \angle DBT$, $\angle BAC = \delta$, $\angle DAC = \gamma$, $\angle BCA = \rho$, $\angle DCA = \mu$, area $ABCD = \triangle$.

Then $\phi = \frac{1}{2}\pi - (\beta - \theta)$, $RB = a\cos(\delta - \theta)$, $RA = a\sin(\delta - \theta)$, $DS = d\cos(\gamma + \theta)$, $AS = d\sin(\gamma + \theta)$, $DQ = \cos(\mu - \theta)$, $CQ = c\sin(\mu - \theta)$, $BP = b\cos(\rho + \theta)$, $PC = b\sin(\rho + \theta)$, $AI + x + CN = AJ + CN = e\cos(\theta)$, $BU + x + DV = BT + DV = f\cos(\theta)$.

 $\therefore BT + DV = f\sin(\beta - \theta)$.

 $x^{2} + \frac{1}{2}x(AI + BU + CN + DV) = \triangle = \frac{1}{2}x(AI + x + CN) + \frac{1}{2}x(BU + x + DV).$

 $\therefore x [e\cos\theta + f\sin(\beta - \theta)] = 2\triangle...(1).$

$$\triangle + \frac{1}{4} \left[a^2 \sin 2(\delta - \theta) + b^2 \sin 2(\rho + \theta) + c^2 \sin 2(\mu - \theta) + d^2 \sin 2(\gamma + \theta) \right. \\ = ef \cos \theta \sin(\beta - \theta) \dots (2).$$

Reducing (2), and remembering that $4 \triangle - 2ef \sin \beta = 0$, we get

$$\tan 2\theta = \frac{a^2\sin 2\delta + b^2\sin 2\rho + c^2\sin 2\mu + d^2\sin 2\gamma - 2ef\sin \beta}{a^2\cos 2\delta - b^2\cos 2\rho + c^2\cos 2\mu - d^2\cos 2\gamma - 2ef\cos \beta}...(3).$$

When a=225, b=153, c=207, d=135, e=240. Then f=277.4, $\triangle=30656.46$, $\delta=38^{\circ}$ 14′ 54″, $\rho=65^{\circ}$ 33′ 40″, $\gamma=59^{\circ}$ 24′ 36″, $\mu=34^{\circ}$ 9′ 22″, $\beta=67^{\circ}$ 3′ 52″.

- $\therefore \tan 2\theta = -.345608 = 2\tan \theta / (1 \tan^2 \theta).$
- \therefore tan θ =5.954833 or -0.167930.
- $\theta = 80^{\circ} 28' 2'' \text{ or } 170^{\circ} 28' 2'' \text{ and } x = -2496.97 \text{ or } -121.044.$

Therefore, two squares can be inscribed in this quadrilateral, the smaller one truly inscribed and the larger with its corners on the sides.

Also solved by J. Scheffer, and V. M. Spunar.

351. Proposed by L. E. DICKSON, Ph. D., The University of Chicago.


Given an isosceles right triangle with hypothenuse h; an isosceles triangle with two sides h and two angles $A{=}22^{\circ}$ 30'; a right angle triangle with the same angle A and opposite side $h/\sqrt{2}$; a triangle with the same angle A, opposite side h, and an angle 45° . Form a triangle whose four pieces are these four triangles, and prove geometrically that it is isosceles.

Solution by G. B. M. ZERR. A. M., Ph. D., Philadelphia, Pa.

Let ABD be the isosceles right triangle with hypothenuse BD=h, sides AB, $AD=h/\sqrt{2}$.

Produce AB to R, making BR=BD and connect RD. Since $\angle ABD=45^{\circ}$, $\angle RBD=135^{\circ}$.

 $\therefore \angle BRD = \angle RDB = 22^{\circ} \ 30' = A$, and BRD is the isosceles triangle with sides h and opposite angles A. Take Q in DA produced so that AQ = AD and join RQ. Then the right triangle ARQ has angle ARQ = A and AQ = h/v/2. Produce AD to P



making DP=h and join RP. Triangle DRP has angle DRP=A, $\angle P=45^{\circ}$, and side DP opposite $\angle DRP=h$.

Now $\angle PQR = 67^{\circ} 30' = 3A$, $\angle ARQ = \angle ARD = \angle DRP = A$.

 $\therefore \angle QRP = 3A$. $\therefore \angle Q = \angle QRP$, and the triangle PQR is isosceles.